

<http://mc.manuscriptcentral.com/jph>

Emerging Public health challenges in the UK: Perception and belief on disproportionate COVID19 deaths among BAME healthcare workforce.

Journal:	<i>Journal of Public Health</i>
Manuscript ID	JPH-20-0482.R3
Manuscript Type:	Short Report
Date Submitted by the Author:	04-Jun-2020
Complete List of Authors:	Moorthy, arumugam; University Hospitals of Leicester NHS Trust, Rheumatology; University of Leicester College of Medicine Biological Sciences and Psychology, Medical Education Sankar, Thangasamy; Kettering General Hospital NHS Foundation Trust
Keywords:	Ethnicity, Health impact assessment, Health services

SCHOLARONE™
Manuscripts

1
2
3 **Emerging Public health challenge in UK: Perception and belief on Increased COVID19**
4 **death among BAME health care workers**
5
6
7
8

9 1. Arumugam Moorthy

10 2. Thangasamy K Sankar

11
12
13 1. Consultant Rheumatologist & Hon Senior Lecturer, Dept of Rheumatology University
14 Hospitals of Leicester NHS trust. Leicester UK.

15 2. Consultant Plastic surgeon, Kettering General Hospitals, Kettering.UK
16
17
18

19 Correspondence author: Dr A Moorthy

20 Email: moorthyarumugam@hotmail.com
21
22

23
24 **Conflicts of interest: None to declare**
25

26 **Funding: None**
27

28
29 **Acknowledgements: Leicester Asian Doctors society and Leicester Asian Nurses society**
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract:**Background:**

Coronavirus infection Disease 19 (COVID19) impacted every part of the world and routine life. Recent report from the Office of national statistics in UK reported disproportionate death among Black Asian and minority ethnic (BAME) population. NHS is heavily relied on the BAME work force both in front line and in the community. We attempted to explore the beliefs and perception about reported worrying issue among BAME health work force in a Diverse city of Leicester.

Methods:

This is a Cross sectional survey using 20 questions in an electronic format. The target population was identified through Leicester Asian Doctors Society and Leicester Asian Nurses Society. The questionnaire was then distributed electronically to the members. Survey questionnaire was accessed by 372, Incomplete response (172) were excluded and 200 completed responses were analysed.

Results:

Majority of BAME workforce are routinely involved in front line duties. More than 70% were anxious about their role during this pandemic. The PPE supply was adequate, and the support received from the local health care providers was more than satisfactory. The work force perceived Comorbidity, Lack of PPE and testing were one of the few reasons for increased death in BAME. BAME group felt adequate provision of PPE, increased testing

1
2
3 and improving mental health wellbeing is required to alleviate concerns and improve BAME
4
5
6 working life in NHS.
7
8
9

10 **Conclusion:**

11
12 BAME workforce are routinely involved in Frontline work and current anxiety level is very
13
14 high. Adequate provision of mental health support with clear risk stratification for return to
15
16 work is required urgently.
17
18
19

20
21
22
23
24
25
26
27
28 Keywords: COVID 19 / Death / BAME group /Health care workers/ BAME health workers
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Background:
4
5
6
7

8 Corona virus infection disease 19 (COVID-19) first case was reported from Wuhan, China in
9 the later part of 2019. COVID-19 rapidly infected other parts of the world, subsequently the
10 World health Organisation declared a Pandemic in March 2020(1). Most of the developed
11 and developing countries went into lockdown measures to avoid morbidity and high
12 mortality rates (2).
13
14
15
16
17
18
19

20 In United Kingdom, COVID19 peaked in April 2020. The Office of national statistics (ONS)
21 data analysed the initial mortality data and reported worryingly disproportionate death
22 among Black, Asian and Minority Ethnicity group (BAME) (3). Ethnicity and disproportionate
23 death rate was reported particularly in Black males are 4.2 times and Black females are 4.3
24 times higher than white ethnicity . However, the adjusted risk for Black Ethnicity is 1.9 for
25 both sexes. This reduction was due to living mainly in London which doubled the risk of
26 anyone living in London. The ONS data also reported people of Bangladeshi, Pakistani,
27 Indian, and Mixed ethnicities also had statistically significant raised risk of death involving
28 COVID-19 compared with those of White ethnicity. Intensive care Audit data reported
29 increase admission of BAME patients in Intensive care unit with COVID-19 (3). It is also
30 reported local authorities such as London and Birmingham with increased proportion of
31 BAME residents and lower income group had increased COVID-19 mortality.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Each percentage point increase in the proportion of the population experiencing income deprivation associated with 2% increase mortality rate [IRR=1.02, 95%CI 1.01–1.04].

1
2
3 UK government swiftly ordered a committee to investigate the reasons for increased death
4 among BAME population and we are now awaiting the report. Several Factors were linked
5 with the increased mortality which includes lower socio-economic status, Social deprivation,
6 Vitamin D Deficiency, Genetics, Co morbid medical conditions and obesity. (11,12).
7
8 Urbanisation is also linked to the disparity in risk and death among BAME workers (28)
9
10
11
12
13
14
15
16
17

18 It is interesting to note the disproportionate death is also high in BAME population in United
19 states (16,). The death rate is high in Black Americans and Asians when compared to white
20 population (17,18). Different fatality rate reported due to COVID-19 in developing countries
21 when compared to developed countries (19,). Various factors linked to increased death in
22 the developing countries are prevalent, particularly co morbidity, poor socio-economic
23 status, and housing, however case fatality rate is low (19). Health inequality and ethnic
24 variation in certain chronic diseases may be another reason and it is one of the Public health
25 challenges currently emerging in UK (20)
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 NHS manpower is hugely relying on international work force particularly from developing
41 countries. Data from NHS (In June 2019) 13.3% of NHS staff in hospitals and community
42 services in England reported a non-British nationality(6).BAME health work force is
43 obviously concerned with disproportionate mortality among BAME health care workers as
44 some of the factors already linked such as Social deprivation, low social economic class,
45 poor housing do not apply. Recent surveys among NHS workforce on COVID including a
46 recent survey by Royal college of Physicians London, among their members also reported
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 48% were very much concerned and this is raised to 76% of those from BAME members
4
5
6 (14).
7
8
9
10
11
12

13 Methodology:

14
15 We conducted a cross sectional survey among BAME health work force in a diverse city,
16
17 Leicester. Leicester hospitals is one of the few trusts employed more BAME Health work
18
19 force in UK. Leicester Hospitals BAME work force comprised Medical and dental staff group
20
21 55.5% and nurses 33.4% among the total local health work force. This may reflect the
22
23 Diverse population of this historic city in the UK. We focused on BAME work force in
24
25 Leicester, a multi culture, diverse city. This is a cross sectional survey among the BAME
26
27 health workforce committed to local health sectors in Leicestershire. We Piloted our initial
28
29 questionnaire, developed and redesigned the questionnaire before circulated the
30
31 anonymised survey electronically. Our questionnaire particularly focused on COVID-19
32
33 among BAME health work force particularly in front line duties and their concerns and
34
35 beliefs. We also attempted to explore the availability of personal protective equipment and
36
37 support received from place of work. Another aspect we explored is about the mental
38
39 health well-being of BAME work force during the COVID-19 pandemic working life. Various
40
41 factors are linked to disproportionate death recently (and we asked the respondents about
42
43 their perception for the possible causes such as co-morbidity, PPE, lack of testing etc.
44
45
46
47
48
49
50
51
52
53

54 The survey was then circulated electronically to BAME health care workers through
55
56 Leicester Asian Doctors society. The Leicester Asian Doctors society (LADS) was established
57
58
59
60

1
2
3 in 2015 by a group of Asian Doctors working in and around Leicester to Support and
4
5 celebrate with a moto of together we care. LADS established a subgroup for Nurses belong
6
7 to BAME group known as Leicester Asian Nurses (LANS). The study period was from 2nd May
8
9 to 17th May, selected few weeks just after the peak of the Epidemic in United Kingdom. The
10
11 study period was selected such a way to get a reasonable immediate reflection from the
12
13 BAME work force following the peak of COVID19 Epidemic in UK. The results were analysed
14
15 using Online software smart survey for statistical purposes.
16
17
18
19
20
21
22
23
24

25 Results

26
27 The response rate for the survey was impressive, with response rate of 60%, when
28
29 compared to our previous survey experience. 374 members accessed our online survey and
30
31 200 completed responses for the analysis. (n=200), 172 incomplete responses were
32
33 excluded. The analysis was carried out using smart survey online software. Our respondents
34
35 are equally distributed with male and female workers in Leicestershire (Chart 1). 78% of
36
37 BAME workers were born outside of the UK and 22 % were born in the UK, with the
38
39 majority, 64 % of workers, being born in India ,11.5% from Black Africans and few from
40
41 Malaysia and Singapore. Our workforce data from our survey reflects the current NHS
42
43 workforce (5,6). The age group ranges from 30 -70 with average age of 40 -50 Years (47%).
44
45 Our respondent split of 70% doctors and 30% from Nursing group. Most of the respondent
46
47 are in full time employment (85.5%) at local NHS trust with University Hospitals of Leicester
48
49 being the main trust. Few respondents are from district general hospital at Kettering and
50
51 Northampton. 30% respondents are from Primary care, 58%from secondary care. In
52
53
54
55
56
57
58
59
60

1
2
3 secondary care there was a spilt of 41% doctors, and 16% nurses. It is interesting to note
4
5 most of the BAME work force (80.5%) are already routinely in front line role. Only 5%
6
7 percentage were deployed to the front line from routine workplace contrary to previous
8
9 survey reported in the media. The exposure to front line work varies from every day 60.1%
10
11 to once every week 11.5%. The current pandemic 66.6% BAME workforce worry, anxious
12
13 about front line which is a cause of concern and this Public health issue was highlighted in
14
15 previous studies (21,22)
16
17
18
19
20
21
22

PPE and Support from NHS:

23
24 Regarding availability and accessibility of PPE 55.5 % had no issues in getting one (Chart 2)
25
26 however, 31.5% had issues with PPE from their workplace. We also asked about any
27
28 difficulty in getting PPE from the managers and noted 66.6% responded that they had good
29
30 support from their senior management team at their workplace (chart 3), and only 16.5%
31
32 need to request or demand PPE at their workplace. 62.5 % of BAME work force are very
33
34 satisfied. Regarding support from Local NHS providers, (chart 5) only 27.5% were not
35
36 satisfied with support from the employers. However, 70.5% respondents are satisfied with
37
38 the trust. 80.5 % respondents were satisfied with the overall support received during the
39
40 pandemic and were satisfied with the support received from the college and national
41
42 society. This demonstrates how best the UK NHS workforce collectively geared up to face
43
44 the unprecedented public health medical emergency in UK.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 *Mental health well-being:*
4

5 Previous studies highlighted the healthcare workers faced anxiety and other mental health
6 issues during COVID19 Pandemic (23,24). Recent studies also emphasised the issue with the
7 mental health well-being of workers during the current pandemic (25). Front-line workers
8 are anxious about their role; however, these workers are self-reliant. There may be a
9 cultural component in handling the uncertainty during pandemic among healthcare workers
10 living and working in isolation in a different country. We explored mental health aspects in
11 our survey. It is interesting to note our respondents (Chart 4) reported that this pandemic
12 had an impact with their mental health. 72% had some form of mental health impact, 55%
13 had mild form, worryingly 11% had a bad impact and out of the 11%, 2% took time off work
14 due to the mental health impact in this pandemic. Only 28% reported no change in their
15 mental health well-being. Our study clearly emphasised more work need to be done
16 urgently to protect the BAME group from mental health issues and offer coping strategies at
17 workplace.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39

40 *BAME COVID 19 Death Rate Beliefs:*
41

42 Various factors and studies were linked to the disproportionate death among BAME group,
43 such as pre-existing co-morbidity, vitamin D deficiency, genetics, lack of PPE and obesity.
44 We asked the respondents about their perception and belief for the disproportionate death
45 among BAME in our survey. It is reported that (chart 5) pre-existing co morbidity (67.5%),
46 lack of PPE (58.5 %), lack of testing (46.5%), vitamin D deficiency (45%), obesity (30.5%)
47 socio economic status (29.5%) in our survey.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 We asked in our survey what changes would help to improve the work environments and
4
5 BAME workforce reported Increasing PPE (71%,) Testing (66 %) and most importantly
6
7 providing adequate Mentorship (70.5%) during this Pandemic at workplace needed
8
9
10 urgently.
11
12
13
14
15
16
17
18
19

20 Discussion:

21
22 Our survey identified the fact most of the BAME Health workers are routinely on Frontline
23
24 duties irrespective of COVID19 pandemic period. BAME workers are in frontline work
25
26 therefore increased risk of infection with COVID 19. It is also highlighted in the report from
27
28 Public health England as clear disparity of risk among BAME group with COVID19 (28). These
29
30 factors lead on to increase level of anxiety among BAME of health workers in UK.
31
32
33

34
35 This study also highlighted the increased level of anxiety among BAME Health workers and
36
37 clear impact on the work force mental well-being during this pandemic. Anxiety level may
38
39 also lead on performance issues at workplace in period of uncertainty and unprecedented
40
41 pandemic. We acknowledge the limitation of the study as it is conducted in a diverse city
42
43 not widely across UK. The survey was conducted among BAME health work force study
44
45 group however Asians are predominately responded with limited response rate from Black
46
47 African Health work force.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 Conclusion:
7

8 From our survey it is evident that the clear majority of the BAME Health care workers are on
9 the frontline in their routine work, therefore they are at higher risk of contracting COVID-19.
10

11 It may be one of the reasons for increased death. Our study respondents perceived BAME
12 doctor's disproportionate death may be due to co-morbidity and lack of PPE and testing.
13

14 With limitation in our study, we are unable to draw a conclusion based on this. Majority of
15 the BAME work force is anxious about the working condition and family particularly living in
16 a social isolation during COVID-19 and reported had significant impact in their mental
17 health. There were no studies comparing the difference of mental health impact in NHS
18 health work force with different ethnicity in a pandemic, clear research is needed exploring
19 the mental health issues in BAME workforce. (23,24)
20

21 Lack of PPE and lack of testing is one of the reasons highlighted for increased death in BAME
22 work force in our survey. It is important to have these issues investigated seriously and
23 clear consistent risks stratification guide for BAME workforce is implemented in workplace
24 urgently. The risk stratification needs to be consistent and should be culturally competent
25 for BAME workforce (25). Culturally competent mental health support for BAME health
26 work force need to be organised by the local health care providers to alleviate the anxiety
27 among BAME workforce (26 ,27). NHS five plan is a welcoming move to reduce the risk and
28 improve the working life of BAME work force in NHS. Evaluation of the newly developed risk
29 stratification framework in real life and implement any innovative measures to improve the
30 confidence of BAME Workforce in NHS (23)
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11 Key messages:
12

- 13 1. Disproportionate COVID 19 Death among BAME group is a public health challenge.
- 14
15 2. BAME Health workers Mental well-being is Impacted with high level of anxiety and
16
17 uncertainty.
18
- 19
20 3. Culturally competent Risk stratification and return to work plan is the need for the
21
22 hour.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17 References:
18

- 19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
1. World Health Organization (WHO) Coronavirus Disease 2019 (COVID-19) Situation Report World Health Organization 2020 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200501-covid-19-sitrep.pdf?sfvrsn=742f4a18_2 (Accessed 26th May)
 2. Office for National Statistics. Coronavirus (COVID-19) related deaths by ethnic group, England <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirus-related-deaths-by-ethnic-group-england-and-wales> (accessed May 14, 2020)
 3. Kings' Fund Report. Ethnic minority deaths and Covid-19: what are we to do? 2020. <https://www.kingsfund.org.uk/blog/2020/04/ethnic-minority-deaths-covid-19> (accessed May 23, 2020).
 4. Watkins J. Preventing a covid-19 pandemic. BMJ. 2020;368: doi: <https://doi.org/10.1136/bmj.m810>
 5. GOV.UK, Ethnicity facts and figures: NHS Workforce. Available at <https://www.ethnicity-facts-figures.service.gov.uk/workforce-and-business/workforce-diversity/nhs-workforce/latest#by-ethnicity>. Accessed on 21st May 2020.

- 1
2
3 6. Immigration and the NHS: the evidence. 19 November 2019 Hugh
4 Alderwick , Lucinda Allen [https://www.health.org.uk/news-and-](https://www.health.org.uk/news-and-comment/blogs/immigration-and-the-nhs-the-evidence)
5 [comment/blogs/immigration-and-the-nhs-the-evidence](https://www.health.org.uk/news-and-comment/blogs/immigration-and-the-nhs-the-evidence)
6
7
8
9
- 10 7. Cook, T., E. Kursumovic, and S. Lennane, Exclusive: deaths of NHS staff from
11 covid-19 analysed. Available at [https://www.hsj.co.uk/exclusive-deaths-of-nhs-](https://www.hsj.co.uk/exclusive-deaths-of-nhs-stafffrom-covid-19-analysed/7027471.article)
12 [stafffrom-covid-19-analysed/7027471.article](https://www.hsj.co.uk/exclusive-deaths-of-nhs-stafffrom-covid-19-analysed/7027471.article). Accessed on 21th May 2020.
13
14
15
16
- 17 8. GOV.UK, Ethnicity facts and figures: NHS Workforce.
18 Available at [https://www.ethnicity-facts-figures.service.gov.uk/workforce-and-](https://www.ethnicity-facts-figures.service.gov.uk/workforce-and-business/workforcediversity/nhs-workforce/latest#by-ethnicity)
19 [business/workforcediversity/nhs-workforce/latest#by-ethnicity](https://www.ethnicity-facts-figures.service.gov.uk/workforce-and-business/workforcediversity/nhs-workforce/latest#by-ethnicity).
20
21
22 Accessed 23rd May 2020
23
24
25
- 26 9. Platt L, Warwick R. Are some ethnic groups more vulnerable to COVID-19 than?
27 others? Institute for Fiscal Studies, Nuffield Foundation. 2020
28
29
30
31
- 32 10. Kirby T. Evidence mounts on the disproportionate effect of COVID-19 on ethnic
33 minorities. Lancet Respir Med. 2020.
34
35 [https://www.thelancet.com/pdfs/journals/lanres/PIIS2213-2600\(20\)30228-9.pdf](https://www.thelancet.com/pdfs/journals/lanres/PIIS2213-2600(20)30228-9.pdf)
36
37
38
- 39 11. Khunti K, Singh AK, Pareek M, Hanif W. Is ethnicity linked to incidence or outcomes of
40 covid-19? Bmj. 2020;369:m1548.
41
42
43
44
- 45 12. Bhalal N, Curry G, Martineau AR, Agyemang C, Bhopal R. Sharpening the global focus
46 on ethnicity and race in the time of COVID-19. Lancet. 2020.
47
48
49
- 50 13. Webb Hooper M, Napoles AM, Perez-Stable EJ. COVID-19 and Racial/Ethnic
51 Disparities. Jama. 2020. doi: 10.1001/jama.2020.8598.
52
53
54
- 55 14. RCP Member survey on work force COVID19 concerns report (Accessed 26th May
56 2020)
57
58
59
60

- 1
2
3 15. Kirby T. Evidence mounts on the disproportionate effect of COVID-19 on ethnic
4 minorities. *Lancet Respir Med*. 2020
5
6
7 16. Clement JM. Knowledge and behaviours toward COVID-19 among U.S. residents
8 during the early days of the pandemic. *JMIR Public Health Surveillance* 2020;6(2)
9 e 19161. doi: 10.2196/19161.
10
11
12
13
14 17. Uma V Mahajan, Margaret Larkins-Pettigrew, Racial demographics and COVID19
15 confirmed cases and deaths: a correlational analysis of 2886 US counties, *Journal of*
16 *Public Health*, , fdaa070, <https://doi.org/10.1093/pubmed/fdaa070>
17
18
19
20
21 18. YancyCWCOVID19andAfricanAmericansJAMA2020 [https://jamanetwork.com/journ](https://jamanetwork.com/journals/jama/fullarticle/2764789)
22 [als/jama/fullarticle/2764789](https://jamanetwork.com/journals/jama/fullarticle/2764789)
23
24
25
26
27 19. Arumugam Moorthy ,Shirish Dubey ,Ash Samanta Ade Adebajo Amita Aggarwal
28 Avinash Jain Nibha Jain S. Sam Lim ,Gail S. Kerr ,Kanta Kumar
29 First published: 30 May 2020 *International Journal of Rheumatic Disease*
30 <https://doi.org/10.1111/1756-185X.1388>
31
32
33
34 20. Kumar K, DubeyS, SamanthaA,; Bosworth A, Moorthy: COVID-19 and Ethnicity:
35 Challenges in Rheumatology Accepted May11 2020 *Rheumatology*
36
37
38
39 21. Shanafelt T, Ripp J, Trockel M. Understanding and Addressing Sources of Anxiety
40 Among Health Care Professionals During the COVID-19 Pandemic. *JAMA*. Published
41 online April 07, 2020. doi:10.1001/jama.2020.5893
42
43
44
45
46 22. Lai J, Ma S, Wang Y, et al. Factors Associated with Mental Health Outcomes Among
47 Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw*
48 *Open*. 2020;3(3):e203976. doi:10.1001/jamanetworkopen.2020.3976
49
50
51
52
53 23. Adams JG, Walls RM. Supporting the Health Care Workforce During the COVID-19
54 Global Epidemic. *JAMA*.2020;323(15):1439–1440. doi:10.1001/jama.2020.3972
55
56
57
58
59
60

- 1
2
3 24. Pareek M, Bangash MN, Pareek N, Pan D, Sze S, Minhas JS, et al. Ethnicity and COVID
4 19: an urgent public health research priority. *Lancet*. 20220
5
6
7
8
9
10
11 25. Risk Reduction Framework for NHS Staff at risk of COVID-19 infection
12 [https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-](https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-on-bame-staff-in-the-nhs/)
13 [on-bame-staff-in-the-nhs/](https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-on-bame-staff-in-the-nhs/) Accessed 25th may 2020.
14
15
16
17
18 26. Addressing impact of COVID-19 on BAME staff in the NHS
19 [https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-](https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-on-bame-staff-in-the-nhs/)
20 [on-bame-staff-in-the-nhs/](https://www.england.nhs.uk/coronavirus/workforce/addressing-impact-of-covid-19-on-bame-staff-in-the-nhs/) Accessed 25th may 2020.
21
22
23
24 27. Albott CS, Wozniak JR, McGlinch BP, Wall MH, Gold BS, Vinogradov S. Battle Buddies:
25 Rapid Deployment of a Psychological Resilience Intervention for Healthcare Workers
26 during the COVID-19 Pandemic [published online ahead of print, 2020 Apr
27 24]. *Anesth Analg*. 2020;10.1213
28
29
30
31
32
33
34 28. Disparities in the risk and outcomes of COVID-19: Public health England Report
35 www.gov.uk accessed 2nd June 2020
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review




Figure 1 : Employment status




Figure 2 PPE Availability at work place




Figure: 3 BAME workforce Impact on Mental well being




Figure: 4 Beliefs for the reasons for increased COVID 19 Death




Figure 5
Support at work place

Or Peer Review